

MORPHOLOGY AND DYNAMICS OF ISOLATED IONOSPHERIC IRREGULARITIES AS DEDUCED FROM GLOBAL NETWORK GPS DATA

E.L. Afraimovich, E.I. Astafieva

We investigate an unusual class of medium-scale traveling ionospheric disturbances (MS TIDs) of the nonwave type, isolated ionospheric irregularities (IIIs) that manifest themselves in total electron content (TEC) variations in the form of single aperiodic negative TEC disturbances of a duration of about 10 min (the total electron content spikes, TECS). For the first time, we present the TECS morphology for 170 days of 1998-2001 with a different level of geomagnetic activity and with the number of stations of the global GPS network ranging from 4 to 240. The number of radio paths used in the analysis totals about 850000. The data were obtained using the technology of global detection and monitoring of ionospheric disturbances (GLOBDET, developed at the ISTP SB RAS) of a natural and technogenic origin using measurements of TEC variations from a global network of receivers of the navigation GPS system [5]. It was found that TECS are observed in no more than 1–2 % of the total number of radio paths. We present the results derived from analyzing the dependence of TECS parameters on the latitude, local time, and on the level of geomagnetic activity, as well as of the correlation between TECS and geomagnetic pulsations. The TECS amplitude exceeds at least one order of magnitude the TEC fluctuation intensity under «background» conditions. The III-induced TEC variations are similar in their amplitude, form and duration to the TEC response to shock-acoustic waves (SAW) generated during rocket launchings and earthquakes. However, the III propagation velocity is less than the SAW velocity (800–1000 m/s) and are most likely to correspond to the velocity of background medium-scale acoustic-gravity waves (AGW), on the order of 100–200 m/s. Preliminary results of our analysis suggest that TECS are often observed during the build-up stage of major earthquakes.